U.S.S.N.: 09/500,904 Filed: February 9, 2000 AMENDMENT

## In the Claims

Claims 1-5 (cancelled)

Patrea Pabst

6. (currently) A diagnostic test to predict the risk of developing lupus comprising reagents which can be used to detect levels of antibodies to Epstein-Barr virus, indicators of Epstein-Barr infection of cells, or levels of Epstein-Barr DNA or protein in a patient, wherein the reagents used to detect antibodies to peptides from Epstein-Barr virus are peptides of up to forty amino acids in length comprising an amino acid sequence selected from the group consisting of PPPGRR? (SEQ ID NO:1), GRGRGRGG (SEQ ID NO:2), RGRGREK (SEQ ID GPORRGGDNHGRGRGRGRGRGGGRPG (SEQ 1D NO:98), GGSGSGPRHRDGVRRPQKRP (SEQ ID NO:25), RPQKRPSC (SEQ ID NO:26), QKRPSCIGCKGTHGGTG (SEQ ID NO:27). GTGAGAGARGRGG (SEQ ID NO:99), SGGRGRGG (SEQ ID NO:100), RGGSGGRRGRGR (SEQ ID NO:101), RARGRGRGRGEKRPRS (SEQ ID NO:102), SSSSGSPPRRPPPGR (SEQ ID NO:103), RPPPGRRPFFHPVGEADYFEYHQEG (SEQ ID NO:104), PDVPPGAI (SEQ ID NO:33), PGAIEQGPA (SEQ ID NO:34), GPSTGPRG (SEQ ID NO:105), GQGDGGRRK (SEQ ID NO:37), DGGRRKKGGWFGKHR (SEQ ID NO:38), GKHRGQGGSN (SEQ ID NO:106), GQGGSNPK (SEQ ID NO:107), NPKFENIA (SEQ ID NO:108), RSHVERTT (SEQ ID NO:109), VFVYGGSKT (SEQ ID NO:110), GSKTSLYNL (SEQ ID NO:111), GMAPGPGP (SEQ ID NO:46), PQPGPLRE (SEQ ID NO:47), CNIRVTVC (SEQ ID NO:48), RVTVCSFDDG (SEQ ID NO:49), PPWFPPMVEG (SEQ ID NO:50) or the peptide consisting of GPQRRGGDNHGRGRGRGRGRGGGRPG (SEQ ID NO:98), or antibodies reactive with these peptides, and

control samples from individuals not at risk of developing lupus, and

OMRF 161 CIP 078617/00140

p. 33

Patrea Pabst

U.S.S.N.: 09/500,904 Filed: February 9, 2000 AMENDMENT

means for determining the differences in levels of antibodies to Epstein-Barr virus, indicators of Epstein-Barr infection of cells, or levels of Epstein-Barr DNA or protein in of a patient and control samples to distinguish individuals at higher risk of developing lupus from those at lower risk of developing lupus.

- 7. (currently amended) The diagnostic test of claim 6 wherein the reagents are used in assays selected from the group of assays based upon the relative presence of an antibody, assays based on cellular proliferation, assays based on molecular binding, assays based on cytokine production, assays based on skin reaction, and assays based on cell surface antigen.
- 8. (currently amended) The diagnostic test of claim 6 wherein the reagents used to detect antibodies to peptides from Epstein-Barr virus are selected from the group consisting of PPPGRRP (SEQ ID NO:1), GRGRGRGG (SEQ ID NO:2), RGRGREK (SEQ ID NO:3), GAGAGAGAGAGAGAGAGAGAGA (SEQ ID NO:7), GPQRRGGDNHGRGKGRGRGRGGGRPG (SEQ ID NO:98), GGSGSGPRHRDGVRRPQKRP (SEQ ID NO:25), RPQKRPSC (SEQ ID NO:26), QKRPSCIGCKGTHGGTG (SEQ ID NO:27), GTGAGAGARGRGG (SEQ ID NO:99), SGGRGRGG (SEQ ID NO:100), RGGSGGRRGRGR (SEQ ID NO:101), RARGRGRGRGEKRPRS (SEQ ID NO:102), SSSSGSPPRRPPPGR (SEQ ID NO:103), RPPPGRRPFFHPVGEADYFEYHQEG (SEQ ID NO:104), PDVPPGAI (SEQ ID NO:33), PGAIEQGPA (SEQ ID NO:34), GPSTGPRG (SEQ ID NO:105), GQGDGGRRK (SEQ ID NO:37), DGGRRKKGGWFGKHR (SEQ ID NO:38), GKHRGQGGSN (SEQ ID NO:106), GQGGSNPK (SEQ ID NO:107), NPKFENIA (SEQ ID NO:108), RSHVERTT (SEQ ID NO:109), VFVYGGSKT (SEQ ID NO:110), GSKTSLYNL (SEQ ID NO:111), GMAPGPGP (SEQ ID NO:46), PQPGPLRE (SEQ ID NO:47), CNIRVTVC (SEQ ID NO:48), RVTVCSFDDG (SEQ ID NO:49), and PPWFPPMVEG (SEQ ID NO:50),

706 283 7737

U.S.S.N.: 09/500,904 Filed; February 9, 2000 AMENDMENT

Apr 05 04 11:30p

- 9. (original) The diagnostic test of claim 8 comprising reagents for detection of
- 10. (original) The diagnostic test of claim 6 for testing patients identified with or at risk of developing systemic lupus erythematosus comprising control samples from individuals with systemic lupus crythematosus.

Claims 11-18 (cancelled)

19. (previously amended) A method for determining the likelihood that an individual has lupus induced by Epstein-Barr virus, or is at risk for developing lupus, comprising obtaining a sample from the individual to be tested,

mixing the sample with reagents which can be used to detect levels of antibodies to Epstein-Barr virus, indicators of Epstein-Barr infection of cells, or levels of Epstein-Barr DNA or protein in a patient,

analyzing the sample, and

comparing the analysis of the sample with results obtained with control samples from individuals not at risk of developing lupus to determine if the differences in levels of antibodies to Epstein-Barr virus, indicators of Epstein-Barr infection of cells, or levels of Epstein-Barr DNA or protein in the individual and control samples indicates the individual is at a higher risk of developing lupus than controls who are at lower risk of developing lupus.

20. (previously amended) The method of claim 19 wherein the reagents are used in assays selected from the group of assays based upon the relative presence of an antibody, assays based on cellular proliferation, assays based on molecular binding, assays based on cytokine production, assays based on skin reaction, and assays based on cell surface antigen.

U.S.S.N.: 09/500,904 Filed: February 9, 2000 AMENDMENT

Claims 23-26 (cancelled).